


Infants' Acetaminophen Concentration Change

Companies are changing the concentration of their medications for infants, and there are new dosing directions for these products.



Professional Dosing Information

What you should know:

- There may be a time when both the old and new concentrations of infants' acetaminophen products are in stores and also in patients' homes
- For children under 2 years of age, no dose is provided on the OTC label — caregivers are instructed to call their doctor for dosing instructions

What you can do:

- 1 Clarify the concentration of infants' acetaminophen product a parent/caregiver is using
- 2 Identify the product concentration and dose for your patient's weight/age from the appropriate chart below
- 3 Discuss the infants' acetaminophen concentration change during sick and well visits

Infants' Acetaminophen

Concentrated Drops — "Old" Concentration

80 mg/0.8 mL

Weight (lb)	Age (mos)	Dose (mL)
6-11 lbs	0-3 mos	0.4 mL
12-17 lbs	4-11 mos	0.8 mL
18-23 lbs	12-23 mos	1.2 mL
24-35 lbs	24-36 mos	1.6 mL

Infants' Acetaminophen

Oral Suspension — "New" Concentration


160 mg/5 mL

Weight (lb)	Age (mos)	Dose (mL)
6-11 lbs	0-3 mos	1.25 mL
12-17 lbs	4-11 mos	2.5 mL
18-23 lbs	12-23 mos	3.75 mL
24-35 lbs	24-36 mos	5 mL

Dosing schedule continues to provide approximately 10-15 mg/kg for children in the 10th-90th decile for weight.¹


Infants' Acetaminophen Concentrated Drops

The enclosed dropper is designed to provide 0.4 mL (40 mg) increments



Infants' Acetaminophen Oral Suspension

The enclosed syringe is designed to provide 1.25 mL (40 mg) increments



Remind parents and caregivers:



- Infants' Acetaminophen Concentrated Drops are 3x more concentrated than Acetaminophen Oral Suspension
- Always read and follow the label
- Keep medicines out of the reach of children
- Never give adult medicines to children
- Use only the dosing device that comes with the product

1. Temple AR. Pediatric dosing of acetaminophen. *Pediatr Pharmacol.* (New York)1983;3(3-4):321-327.


Advil®

Relief you can trust

Measure dosing only with the device that comes with the product.



Available in three flavors!
• Fruit • Grape • Blue Raspberry



	Infants' Advil® Concentrated Drops for ages 6-23 months	Children's Advil® Suspension for ages 2-11 years	Advil® Tablets Advil® Liqui-Gels® for ages 12 and up
Active ingredients	Ibuprofen 50 mg (NSAID)* in each 1.25 mL	Ibuprofen 100 mg (NSAID)* in each 5 mL	Ibuprofen 200 mg (NSAID)* in each tablet
Indications	Fever reducer, relieves minor aches and pains due to cold, flu, headaches, and toothaches	Fever reducer, relieves minor aches and pains due to cold, flu, sore throat, headaches, and toothaches	Fever reducer, relieves minor aches and pains due to headache, toothache, backache, menstrual cramps, cold, muscular aches, minor pain of arthritis

		Weight	Age				
		under 6 months		do not use	do not use	do not use	
Patient Dosing	12-17 pounds	6-11 months	1.25 mL[†] every 6-8 hrs not to exceed 4 doses/24 hrs	↓	Use Infants' Advil® Concentrated Drops	Use Infants' Advil® Concentrated Drops	
	18-23 pounds	12-23 months	1.875 mL[†] every 6-8 hrs not to exceed 4 doses/24 hrs				
	24-35 pounds	2-3 years	Use Children's Advil® Suspension	↓	1 tsp [†] every 6-8 hrs not to exceed 4 doses/24 hrs	Use Children's Advil® Suspension	
	36-47 pounds	4-5 years					1½ tsp[†] every 6-8 hrs not to exceed 4 doses/24 hrs
	48-59 pounds	6-8 years					2 tsp[†] every 6-8 hrs not to exceed 4 doses/24 hrs
	60-71 pounds	9-10 years					2½ tsp[†] every 6-8 hrs not to exceed 4 doses/24 hrs
	72-95 pounds	11 years					3 tsp[†] every 6-8 hrs not to exceed 4 doses/24 hrs
		12+ years	Use Advil® Tablets or Liqui-Gels®	Use Advil® Tablets or Liqui-Gels®	1 or 2 tablets[†] every 4-6 hrs not to exceed 6 tablets/24 hrs		

*Nonsteroidal anti-inflammatory drug.

[†]Measure dosing only with the device provided.

[†]If pain or fever does not respond to 1 tablet, 2 tablets may be used.

Please refer to drug facts on the product label for complete warnings.

Important Instructions for Proper Use

- Read and keep the cartons for complete warnings and information on Advil® products
- Remind patients that if they are taking other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), they should consult with you before taking any Advil® products
- Measure dosing only with the device provided
- Adult strength medicines should not be administered to children
- Products contain other ingredients. Please see full ingredient list on each package

Visit advil.com for drug facts, product updates, and additional information.